



Available online at www.sciencedirect.com

ScienceDirect



REVIEW

Yield gap and resource utilization efficiency of three major food crops in the world — A review

RONG Liang-bing^{1,2,3}, GONG Kai-yuan^{1,2,3}, DUAN Feng-ying³, LI Shao-kun³, ZHAO Ming³, HE Jian-qiang^{2,4,5}, ZHOU Wen-bin³, YU Qiang^{2,6,7}

¹ College of Natural Resources and Environment, Northwest A&F University, Yangling 712100, P.R.China

² State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, Institute of Soil and Water Conservation, Northwest A&F University, Yangling 712100, P.R.China

³ Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing 100081, P.R.China

⁴ Key Laboratory for Agricultural Soil and Water Engineering in Arid Area, Ministry of Education, Northwest A&F University, Yangling 712100, P.R.China

⁵ Institute of Water-Saving Agriculture in Arid Areas of China, Northwest A&F University, Yangling 712100, P.R.China

⁶ School of Life Sciences, University of Technology Sydney, Broadway, NSW 2007, Australia

⁷ College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, P.R.China

Abstract

Yield gap analysis could provide management suggestions to increase crop yields, while the understandings of resource utilization efficiency could help judge the rationality of the management. Based on more than 110 published papers and data from Food and Agriculture Organization (FAO, www.fao.org/faostat) and the Global Yield Gap and Water Productivity Atlas (www.yieldgap.org), this study summarized the concept, quantitative method of yield gap, yield-limiting factors, and resource utilization efficiency of the three major food crops (wheat, maize and rice). Currently, global potential yields of wheat, maize and rice were 7.7, 10.4 and 8.5 t ha⁻¹, respectively. However, actual yields of wheat, maize and rice were just 4.1, 5.5 and 4.0 t ha⁻¹, respectively. Climate, nutrients, moisture, crop varieties, planting dates, and socioeconomic conditions are the most mentioned yield-limiting factors. In terms of resource utilization, nitrogen utilization, water utilization, and radiation utilization efficiencies are still not optimal, and this review has summarized the main improvement measures. The current research focuses on quantitative potential yield and yield gap, with a rough explanation of yield-limiting factors. Subsequent research should use remote sensing data to improve the accuracy of the regional scale and use machine learning to quantify the role of yield-limiting factors in yield gaps and the impact of change crop management on resource utilization efficiency, so as to propose reasonable and effective measures to close yield gaps.

Keywords: food crops, yield, yield gap, resource utilization efficiency, yield-limiting factor

Received 14 September, 2020 Accepted 16 December, 2020
RONG Liang-bing, E-mail: rongliangbing@163.com;
Correspondence ZHOU Wen-bin, E-mail: zhouwenbin@caas.cn;
YU Qiang, E-mail: yuq@nwfau.edu.cn

© 2021 CAAS. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
doi: 10.1016/S2095-3119(20)63555-9

1. Introduction

The ability to feed the world's growing population depends on the ability of future food supplies to meet food demand (Cirera and Masset 2010). However, the current rate of grain

yield increases suggests that there will be a crisis involving insufficient grain supply in the future. The United Nations predicts that the population of the earth will increase by 9.7 billion in the next 30 years (UNDESAPD 2019). In addition, changes in the structures of diets for people across the globe have further increased the demand for crops. Calorie consumption has increased from 2 250 kcal per capita per day in the 1860s to 2 880 kcal in 2015, and it is estimated that it will reach 3 900 kcal by 2050 (Pardey *et al.* 2014; Valin *et al.* 2014), leading not only to increased demand for food crops, but also to significantly increased demand for crops to feed livestock. Therefore, FAO believes that ensuring food security will require global crop yields to increase by more than 70% (FAO 2009).

There are two main ways to increase grain yield: one way is to increase the area of cultivated land, and the other way is to increase the yield of existing cultivated land. Many studies have shown that there is a large amount of land suitable for crop growth in northern South America and tropical Africa that could be converted into cultivated land (Rosegrant *et al.* 2001; Bruinsma 2003). However, most of these land is located in tropical rain forests or nature reserves, both of which have high social, economic and ecological values. Additionally, loss of forest cover in tropical soils results in rapid loss of soil fertility, requiring additional investments to preserve soil organic matter (Ramankutty *et al.* 2002; Licker *et al.* 2010). For many countries, the area of arable land that can be increased is limited, and the existing arable land area is gradually decreasing due to land degradation, urbanization and resource extraction (Lu *et al.* 2007; Fitzgerald *et al.* 2020). In fact, the agricultural land area in developed countries decreased by 34% from 1995 to 2007, and the cultivated land area in developing countries increased by 17.1%, mainly in tropical forests (Gibbs *et al.* 2010). Therefore, it is difficult to achieve a substantial increase in global food production by increasing the area of cultivated land.

The green revolution in the second half of the 20th century greatly increased the yield of farmland. Global crop yield tripled from 1960 to 2014 at the same time that arable land increased by only 10% (Pellegrini and Fernández 2018). These developments have led researchers and policy makers to investigate the prospect of meeting food demand through increased yields. However, some studies have pointed out that the current crop yield growth rate in major food-producing countries has slowed or even stalled. Since the 1990s, for instance, rice (*Oryza sativa* L.) yields in Asia have hardly increased at all (Cassman *et al.* 2003).

We have collected and summarized the research on the yield gaps and resource utilization efficiency (RUE) gaps of the world's major food crops in recent years. Our analysis of the yield gaps and RUE of different crops in different

countries will provide some guidance for increasing crop yields in the future.

2. Yield gap

The study of yield gap began in the 1970s with the International Rice Research Institute's study of yield-limiting factors for rice in six Asian countries (Inst *et al.* 1979). The concept of yield gap was first proposed in 1981, and was defined as the gap between actual farm yield and experiment station yield (de Datta 1981). de Datta (1981) also defined potential farm yield to represent the highest yield that farmers could achieve. Based on actual farm yield and experiment station yield, "technical upper limit yield" and "economic upper limit yield" were introduced (Fresco 1984). The technical upper limit yield is the maximum yield that a specific piece of farmland can reach, which is equivalent to the potential farm yield. The economic upper limit yield refers to the yield obtained when the farm profit is the highest. Because the input level used by farmers is lower than the input level required for the maximum yield for a specific piece of farmland, the economic upper limit yield is lower than the technical upper limit yield. Some researchers have subsequently defined "potential yield" and "available yield" (Rabbinge 1993). Potential yield is the yield that could possibly be achieved under the cultivation and management practices used at an experimental station to ensure good crop growth (i.e., using suitable varieties, weed control, pest management, optimum planting dates and seeding rates, not subject to water and fertilizer stress, limiting harvest losses, etc.). The potential yield reflects the local yield that is determined only by light and temperature conditions and genetic characteristics. Available yield is the yield that can actually be achieved under optimal cultivation management measures, and it is almost the same as the potential farm yield and the technical upper limit yield. It is now generally accepted that 80% of the potential yield should be considered as the attainable yield (Cassman *et al.* 2003; Lobell *et al.* 2009). de Bie (2000) summarized previous research on yield gap, and identified several yield levels: the potential yield of an experimental station simulated by a model, the maximum yield of the experimental station, farmers' potential yield, the economic upper limit yield, and the actual farmland yield. Lobell and Ortiz-Monasterio (2006) defined "field yield gap" as the difference between the highest field yield and the regional average yield.

Previous yield gap research has usually divided yield into different levels, thus resulting in different definitions of the yield gap. In addition to the yield levels mentioned above, the two yield levels of "basic soil fertility" and "high-yield and high-efficiency" also have research value. However, there

is currently almost no relevant literature related to these two yield levels. “Basic soil fertility” refers to the crop yield level obtained without the input of irrigation water and applied fertilizer, relying only on the soil’s native fertility and existing climatic conditions. It represents the local minimum yield. “High-yield and high-efficiency” is a concept proposed by Chinese researchers. Designed management practices consist of a crop module from which cropping strategies in a given region and a resource supply module for the formulation of nutrient and water applications according to soil tests and the needs of the growing crops. It represents the application of comprehensive and integrated agricultural technology innovations to achieve high crop yields and efficient utilization of resources (Shen *et al.* 2013; Wang *et al.* 2014; Bai *et al.* 2020), and is the same as the “Nutrition Expert System” described by Xu *et al.* (2016). The farm yield is the actual yield achieved by farmers. The gap between “basic soil fertility” yield and farm yield can reflect the benefits of using agricultural resource inputs, and is helpful for further research on evaluating RUE. The gap between farm yield and “high-yield and high-efficiency” yield reflects the increase in farm yield when resource utilization is optimal. Fig. 1 is an expansion of the yield gap model presented by de Bie (2000) that now includes “high-yield and high-efficiency” yield and “basic soil fertility” yield.

2.1. Quantitative methods for yield levels

Different yield levels are measured in different ways. Farm

and experimental station yield levels can be obtained through household surveys and field trials. Field trials can control factors affecting yields by imposing different treatments in order to analyze the impact of specific factors on crop yield formation. Meertens (1991) applied six treatments that included weeds, fertilizers, pests, and diseases, and found that only weed treatments had a significant effect on cotton yield. However, the results of the field test only reflected the local situation during the current year of the study, and may not be applicable to regional scales and long-term sequences. Farm household surveys can estimate different yield levels and yield gaps at the regional scale by obtaining information such as soil, crop varieties and farm household management. Sarobol *et al.* (1989) used household surveys and related statistical data, and found that the limiting factors for soybean yield in Thailand were mainly unreasonable land use, weeds, insect pests, inadequate land preparation, and outdated production technologies. However, household surveys usually use questionnaires that are subjective and may omit relevant information about the causes of yield gap. These types of questionnaires make it difficult to obtain accurate farmland management information.

Since the potential yield cannot be obtained through actual production data, its quantification is usually achieved by use of simulation models. The models used to determine yield gaps can be divided into two categories: empirical models and mechanistic models (Cao *et al.* 2011). Empirical models are mathematical statistical analysis models based on the statistical correlation between biomass and climate




Fig. 1 Yield gaps and their main constraints. Yield Gap-0 cannot be reduced; Yield Gap-1 is the gap between the yield produced at experimental station level and farmland level; Yield Gap-2 is the gap caused by different management practices at the farmland level; Yield Gap-3 is the gap between farmland and wasteland, and caused by water, fertilizer, and other management measures. This figure is an expansion of the figure presented by de Bie (2000).

factors, such as the Miami Model (Luo *et al.* 2011; Cheng *et al.* 2012; Yu *et al.* 2012), the Thornthwaite Memorial Model (Mao *et al.* 2008; Li *et al.* 2009), the Wagenigen Model (Lin *et al.* 2003), and the AEZ Model (Liu *et al.* 2001). Mechanistic models use mathematical concepts to simulate crop physiological processes and to explain the overall function of the crop, such as WOFOST (Wu *et al.* 2006), Hybrid-Maize (Liu *et al.* 2017), DSSAT (Maria Carolina *et al.* 2018), EPIC (Lu and Fan 2013), APSIM (Li *et al.* 2014), ORYZA (Agus *et al.* 2019), etc. In addition to simulating potential yield, a mechanistic model can also simulate other yield levels by defining different cultivation scenarios (Bindraban *et al.* 2000). Because these models are often built on the basis of single-point experiments, there may be many limitations in using them to calculate regional yield gaps. Regional yield gap studies have often used the method of combining models and remote sensing (Maas 1988; Moulin *et al.* 1998). For example, Li *et al.* (2014) combined the APSIM-Wheat Model with GIS technology to analyze the temporal and spatial distribution of the winter wheat yield gap in the North China Plain.

It should be noted that using different methods to estimate potential yield (Y_p) has a significant impact on the calculation of the yield gap. Taking the North China Plain as an example, the relative yield calculated using a crop model was low (56–68%) (Lu and Fan 2013; Li *et al.* 2014; Fang *et al.* 2017). However, the relative yield value calculated using boundary line analysis was 85% (Cao *et al.* 2019). This difference in relative yield is caused by the difference in the definition of potential yield. The Y_p obtained by the crop model is the theoretical maximum yield, while the potential yield estimated based on field trials is the yield that high-yield farmers can actually achieve.

2.2. Current status of world grain crop yield

At present, the primary focus of yield gap research is centered around the three major food crops of wheat (*Triticum aestivum* L.), maize (*Zea mays* L.) and rice. Crop yield data for 2002–2017 were obtained from Food and Agriculture Organization Statistics (www.fao.org/faostat, accessed on 20 June, 2020), maize yield was the highest at 4.3 t ha⁻¹, followed by rice (3.7 t ha⁻¹) and wheat (3.1 t ha⁻¹). Ireland had the highest wheat yield, and western Europe was a high-yield area for wheat, almost countries in this region could produce more than 6 t ha⁻¹. High-yield areas for maize were found on the Arabian Peninsula, including the United Arab Emirates (UAE), Israel, Jordan, and Kuwait, maize yields where were more than 18 t ha⁻¹, while most countries had less than 10 t ha⁻¹. There was no single high-yield area for rice, but rice yield in Egypt and Australia exceeded 9 t ha⁻¹. And for all three crops, the countries with low yields

were concentrated in Africa (Appendix A).

Yield gap data were obtained from the Global Yield Gap and Water Productivity Atlas (Appendix B) (www.yieldgap.org accessed on 30 April, 2020) and other literature (Appendix C), including actual yield (Y_a) and Y_p for countries or regions. Relative yield was calculated as:

$$\text{Relative yield} = Y_a / Y_p \quad (1)$$

where Y_p values for wheat, maize and rice were 7.7, 10.4 and 8.5 t ha⁻¹, respectively. In general, wheat and maize production in Europe, especially in western Europe, is close to potential production; the relative yields of the three crops in Africa are low, and many countries have less than 40% (Lobell *et al.* 2009; Hoffmann *et al.* 2018); the relative yields of America and Asia are similar, higher than that in Africa but lower than that in Europe; the relative maize yield of Australia exceeds 80%, but wheat is only 48%. In particular, the relative yields of maize and rice in the United States are between 60–80%, but the yield of wheat can only reach 30% of the potential yield, which may be caused by lack of poor soil quality (Patrignani *et al.* 2014); while relative yield of rice in Egypt is the highest among all countries with data records, benefiting from Egypt's policies reforms, technological change and increased investment (Fan *et al.* 1997). However, the harvest in Africa is only 20% of potential yield (Appendices B and C).

For each of the three major crops, we have focused on the analysis of the yield gap in four major producer countries, two low-yield-gap countries and two high-yield-gap countries. Y_{a1} , Y_{a2} and Y_{a3} are the abbreviations for the average actual yields from 1961 to 1980, 1981–2000, and 2001–2018, respectively.

The yield of wheat continued to increase from 1961 to 2018, but the gap between Y_{a3} and Y_{a2} in all eight countries was smaller than the gap between Y_{a2} and Y_{a1} (Fig. 2). These results demonstrate a declining rate of wheat yield increases over time. If 80% of Y_p is used as attainable yield, Ethiopia will aim to increase yield by 4.6 t ha⁻¹, the largest increase of the eight countries. Tanzania would aim to increase production by 240%, high-yield-gap countries have the greatest room for increase yield. In contrast, it is continue to increase yield in Denmark and Germany almost impossible to due to the current yield's closeness to Y_p . For the four major wheat-producing countries, although the increase in yield has declined, it has not yet reached the threshold of attainable yield.

Since the beginning of the 21st century, maize yields have continued to increase substantially, while in China and Nigeria, the growth trends have been weakening (Fig. 3). In the USA, Germany and Netherlands, current relative yield close (or exceed) to 80%. Compared with increasing yield, these countries need to maintain the stability of production and reduce the inter-annual fluctuation of yield




Fig. 2 Actual yield (Ya) and potential yield (Yp) of wheat in four major wheat-producing countries (China, France, India, and USA), low-yield-gap countries (Denmark and Germany), and high-yield-gap countries (Ethiopia and Tanzania). Ya1, Ya2 and Ya3 are average Ya during 1961–1980, 1981–2000 and 2001–2018, respectively. Bars mean SD ($n=20$).




Fig. 3 Actual yield (Ya) and potential yield (Yp) of maize in four major maize-producing countries (Argentina, Brazil, China, and USA), low-yield-gap countries (Germany and Netherlands) and high-yield-gap countries (India and Nigeria). Ya1, Ya2 and Ya3 are average Ya during 1961–1980, 1981–2000 and 2001–2018, respectively. Bars mean SD ($n=20$).

(SD in Fig. 3). The actual yield of India and Nigeria is less than 20% of the potential yield, and there is great potential for increasing yield through changes in crop management in the future. Actually, in Africa, backward farming and management practices, limitations on use of water and fertilizers, and pest and disease pressures have led to low maize yields for farmers (van Ittersum *et al.* 2013; van Loon *et al.* 2019). Therefore, introduction of advanced technology for managing maize production in these areas is urgently needed.

Rice production is mainly concentrated in Asia. As the largest rice producer, the relative yield of rice in China is close to 70%, with a stable yield and little increase in the most recent 20-yr period (Fig. 4). Late rice in the middle and lower reaches of the Yangtze River has even reached 90% of the potential yield, with almost no increases in yield in recent years (Zhang *et al.* 2019b). Bangladesh, India, Cote D'Ivoire and Zambia have still maintained relatively higher growth rates than China, but the relative yield of less than 50% shows that rice yield still has potential to increase rapidly with technical support. Egypt is special, although its relative yield has reached 81%, its rice yield still maintains a high growth rate.

In general, the greater the gap between Yp and Ya, or the smaller the relative yield, the easier it is to increase crop yield. Therefore, low-yield areas should be given priority over high-yield areas in developing production methods to narrow the yield gap (Wilbois and Schmidt 2019). Farm yield in developed countries and regions such as the United States and Europe is close to being at the threshold of 80% of potential yield. Although it can be further improved by technical means, the return on investment for increased yield




Fig. 4 Actual yield (Ya) and potential yield (Yp) of rice in four major rice-producing countries (Bangladesh, China, India, and Indonesia), low-yield-gap countries (China and Egypt), and high yield gap countries (Cote D'Ivoire and Zambia). Ya1, Ya2 and Ya3 are average Ya during 1961–1980, 1981–2000 and 2001–2018, respectively. Bars mean SD ($n=20$).

is very low. In contrast, the relative yield of farmers in Africa and South America is low, and these areas are the focus of reducing the yield gap in the future, as confirmed by the prioritization provided by van Oort *et al.* (2017).

2.3. Factors causing yield gap and ways to close the gap

Factors causing yield gaps are called yield-limiting factors. There are many of these factors, such as soil quality, genetic

factors and human management (including irrigation, fertilization, pest management, and planting factors) (Licker *et al.* 2010). Usually the yield gap is directly used to quantify yield-limiting factors. For example, the water limitation can be quantified by determining the yield gap between rainfed and irrigated yields, and the nutrient limitation can be quantified by determining the yield gap between different fertilizer application amounts. In addition, many studies have adopted the boundary line analysis method (Fermont *et al.* 2009; Hajjarpoor *et al.* 2018; Cao *et al.* 2019). The upper boundary line is used to minimize the influence of other factors, and the response function of yield to the limiting factor is obtained. Compared with the previous method, it can get a continuous result that a certain factor limits the yield. According to the above research (Appendix C), the main factors limiting crop production are shown in Fig. 5 and Appendix D.

Climate Changes in atmospheric CO₂, temperature and precipitation are the main driving forces affecting crop response to climate change (Hatfield *et al.* 2011). Many studies have shown that climate warming will reduce crop yields to a certain extent, and climate warming results in positive yield responses in only a few regions such as Northeast China (Tao *et al.* 2006; Liu Y *et al.* 2010; Asseng *et al.* 2013; Kukul and Irmak 2018). This may be due to the high latitude of these areas, and current temperatures not being in the range to promote maximum growth and yield. Increasing atmospheric CO₂ concentrations promote CO₂ fertilization effects on plants, but the amount of yield that can

be increased by increased CO₂ has not yet been determined (Long *et al.* 2006; Tao *et al.* 2006; Ewert *et al.* 2007). The regional trend of rainfall is obvious. Precipitation in arid areas of China has decreased in summer and autumn. In contrast, in the humid southern areas of China, there has been more precipitation in summer, negatively impacting agricultural production (Fan *et al.* 2011). In addition, studies on rice production in Southeast Asia have shown that there is a significant difference between yields in the rainy season and the dry season. The relative rice production in Indonesia in the dry season is only 55%, but it can reach 67% in the rainy season. However, the opposite is true in Vietnam (75% in the dry season and 50% in the rainy season) (Laborte *et al.* 2012; Silva *et al.* 2017).

Nutrient and water management The growth and development of crops are inseparably linked to water and nutrient availability. Generally, the higher the input level is, the higher the yield is. Among the many inputs used in agricultural production, water and nitrogen are the most important. 10.7% of the single-season rice yield gap in Northeast China is due to insufficient nitrogen fertilizer supply, while 7.2% is due to water stress (Zhang *et al.* 2019b). Also, there is a connection between water and nitrogen. In the western United States, nitrogen fertilizer accounted for 24% of the yield increase in a maize-soybean rotation under irrigation. However, without irrigation, the contribution of nitrogen fertilizer could be ignored (Balboa *et al.* 2019). In addition to nitrogen, other nutrients (such as phosphorus, potassium, etc.) will also limit crop yields (Hajjarpoor *et al.* 2018). In particular, the lack of potassium fertilizer produced a greater yield restriction on maize production than that on rice and wheat production (Jin 2012; Dai *et al.* 2013; Xu *et al.* 2016).

Varieties The increase in crop yields since the Green Revolution is related to the selection of new varieties. In the early Green Revolution (1961–1980), new varieties accounted for 17% of the yield growth in developing countries. However, by the late Green Revolution period (1980–2000), this proportion had increased to 50% (Evenson and Gollin 2003). Senapati and Semenov (2020) believed that breeding varieties that are resistant to heat and drought at the flowering stage, and have the best canopy and root structure could increase wheat yield by 3.5–5.2 t ha⁻¹. Liu *et al.* (2012) found that new maize varieties increased maize yield by extending the growing season. In China, more than half of the farmers did not choose the right corn varieties, which caused a 19.8% yield gap (Zhang *et al.* 2016). Improved varieties have also played an important role in increasing rice yield (Takai *et al.* 2006; Peng *et al.* 2008; Espe *et al.* 2018), and the best varieties should have longer growing seasons, larger spikes and greater kernel weight (Zhang *et al.* 2019b). In order to obtain higher yields




Fig. 5 The frequency of the main yield-limiting factors in the published papers, including climate, nutrient, water, varieties, planting date (or seeding date), and socioeconomic factors. Socioeconomic factors include agro-technical service, government support, risk aversion of farmers, farm size, labor, and expertise, etc.

in the United States under the pressures of climate change, pests and diseases, crop production systems widely use hybrids and herbicide-resistant varieties that have better adaptability to the environment (Nalley *et al.* 2016).

Planting date Temperature, light, precipitation, and solar radiation play an important role in crop growth and development, but it is difficult for humans to directly control these parameters in field production situations. By adjusting the planting date, crops can, to a certain extent, make better use of these resources. Khaliq *et al.* (2019) used the APSIM Model to simulate the effect of planting date on rice and wheat yields. They found that the best planting date for rice in Pakistan was one month later than the currently used planting date, and that the best planting date for wheat was at least one month earlier than the currently used planting date. The suitable planting period for winter wheat in the North China Plain is from October 3 to October 8, and the planting period is the largest limiting factor affecting the number of heads (Cao *et al.* 2019). The transplanting date for rice will affect the number of days in the growing season and the climatic conditions received during the growing season (Hu *et al.* 2017). The transplanting date for rice in Northeast China has a greater impact on yield than it does in the middle and lower reaches of the Yangtze River (Zhang *et al.* 2019b). Many studies have shown that temperate regions facing the negative impacts of global warming can increase production through earlier planting and emergence. Some of these studies have involved wheat in Australia (Zeke and Nendel 2016) and rice in China and the United States (Hardke *et al.* 2013; Linquist and Espe 2015; Ding *et al.* 2020). In areas with sufficient heat, the planting date needs to be properly delayed to avoid damage to crops caused by high temperature (Ding *et al.* 2020).

Socioeconomic factors The goal of farmers involved in agricultural production is generally to obtain maximum economic benefits rather than to obtain the highest yield. Reducing yield gap through better management measures may not be feasible in all regions because it means higher economic and resource investments. Even though the yield of improved varieties in Africa can be three times greater than the average yield obtained by farmers, they often do not consider using these improved varieties due to the higher seed costs and drought intolerance (van Loon *et al.* 2019). In Northwest China, farmers have been unable to increase yield through large-scale wheat and maize intercropping due to water shortages because both intercropping and increased planting density result in higher water demand (Fang *et al.* 2017). In addition, factors such as the size of the farm (Dutta *et al.* 2020), the number and gender of the labor force (Mahmood *et al.* 2019) and the education level of farmers (Assefa *et al.* 2020) can affect the decision-making processes of farmers, and ultimately the final yields

obtained.

Other yield-limiting factors affect yield gaps less, but this does not mean that those factors are unimportant. Liu W *et al.* (2010) established a planting density gradient experiment and found that maize yield increased by 48–72% under high plant density conditions. Flat land allows soil moisture and nutrients to be evenly distributed, and also results in production areas that are conducive to the use of agricultural machinery to manage farmland in an efficient manner to increase yields. In the continuously cropped rice production areas of California (USA), precision land-leveling began to be promoted as early as the 1970s (Espe *et al.* 2016). Altitude is generally ignored in field trials, but when Wang *et al.* (2020) used remote sensing to analyze differences in rice yield in Northeast China, they found that, on the regional scale, altitude was one of the important limiting factors for yield. Each 1-m increase in altitude resulted in a potential yield reduction of 8.6 kg ha⁻¹.

3. Resource utilization efficiency

Even though crop yield is often limited by factors such as light, temperature, water, and nutrients, current yield gap research has focused on improving crop yields without considering RUE. In fact, even though excessive inputs of resources can increase yields, the utilization efficiency of those resources will decrease. This approach of applying excessive amounts of inputs not only reduces the economic benefits received by farmers, but also causes environmental pollution. Therefore, it is necessary to consider RUE when studying how to narrow yield gap. Research on efficiency gap is currently focused on nutrients, water and radiation. This type of research is generally conducted by setting up different management practices, and then comparing RUE under each practice.

3.1. Nutrient utilization efficiency

There are many indicators for evaluating nutrient utilization efficiency. Commonly used indicators are agronomic efficiency (AE), recovery efficiency (RE) and partial factor productivity (PFP). Taking nitrogen fertilizer utilization efficiency as an example, the calculation equations are:

$$NAE=(Y-Y_0)/F \quad (2)$$

$$NRE=(U-U_0)/F \quad (3)$$

$$NPFP=Y/F \quad (4)$$

where NAE is agronomic efficiency of nitrogen fertilizer, Y is grain yield when nitrogen fertilizer is applied, Y₀ is grain yield without nitrogen fertilizer, F is nitrogen fertilizer application amount, NRE is nitrogen fertilizer recovery efficiency, U is the amount of nitrogen absorbed by the crop when nitrogen fertilizer is applied, U₀ is the amount of nitrogen absorbed

by crops without nitrogen fertilizer, and NPFP is nitrogen partial factor productivity.

AE directly reflects the impact of fertilizer application on production and is related to economic returns; RE focuses on the nutrient response of crops, and like AE, it requires plots without nutrient input as a control, and requires long-term trials to eliminate residual effects; PFP is a simple expression of factors of production. Compared with AE and RE, the data required for PFP is easier to obtain (Fixen *et al.* 2015) (Fixen *et al.* 2015). NPFP values of three crops in different countries are shown in Appendix A.

Among the three crops evaluated in this review paper, the current yield of maize is closer to Y_p than the yields of wheat and rice, and therefore agronomic management of maize is likely being conducted in a more reasonable and efficient manner. Analysis of NPFP from 2002 to 2017 (www.fao.org/faostat, accessed on 20 June, 2020) in the major maize-producing countries (United States, China, Brazil, and Argentina) and advanced production countries (Appendices B and C with relative yield greater than 80%) showed that as fertilizer input increases, nutrient utilization efficiency decreases (Fig. 6), even though yield increases as fertilizer input increases. Therefore, the methods used to increase the amount of fertilizer applied in the studies in order to reduce the maize yield gap were not conducive to the efficient utilization of nutrients. In fact, the process of nutrient absorption is affected by more than just the amount of nutrient input. Nutrient absorption is also the result of a combination of factors. Despite the same level of nitrogen input, NPFP values for maize in China were lower than those observed in the advanced production countries. This result shows that in China and Brazil, it is still possible to improve nitrogen use efficiency and production of maize by reducing the negative impacts of other yield-limiting factors. The relative yield of the United States was close to 80%, and its NPFP had reached the level of advanced producing countries. Argentina's NPFP was somewhat higher than the ideal condition defined by the regression relationship in Fig. 6, which may be a result of its fertile soil and suitable climate.

Some studies have shown that excessive fertilization would lead to low nutrient utilization efficiency (Mueller *et al.* 2012; Cui *et al.* 2014; Chen *et al.* 2018). Increasing irrigation amounts and planting density are common ways to improve nitrogen fertilizer utilization efficiency (Zhang *et al.* 2019a; Bai *et al.* 2020). Changing the fertilization formula (Li *et al.* 2017), cereal/legume intercropping (Xu *et al.* 2020) and fertilization time (Meng *et al.* 2016) can also improve the efficiency to a certain extent. In addition, some studies have shown that a co-limiting relationship exists between the absorption of different nutrients, i.e., the lack of one element will lead to a reduction in the absorption of




Fig. 6 Relationship between nitrogen rate and nitrogen partial factor productivity (NPFP) for maize. Advanced production countries (relative yield exceeding 80%) include Australia, Belgium, Germany, Netherlands, and Italy.

another element. Therefore, nutrient utilization efficiency can be improved by adjusting the ratio of nitrogen-sulfur or nitrogen-phosphorus (Sadras 2006; Carciocchi *et al.* 2020). In the maize production system used in northeastern China, if the farmers adjust the typical fertilization amounts (207 kg N ha^{-1} , 40 kg K ha^{-1} and 61 kg P ha^{-1}) to the fertilizer amounts recommended by nutrition experts (173 kg N ha^{-1} , 32 kg K ha^{-1} and 69 kg P ha^{-1}), the partial productivity of nitrogen and phosphorus will increase by 24.6 and 12.6%, respectively, and maize yield will increase by 0.9 t ha^{-1} (Xu *et al.* 2016).

3.2. Water utilization efficiency

Water utilization efficiency (WUE) is an important indicator to measure the relationship between crop yield and water consumption. However, due to different research objectives, the calculation method for WUE can be different. For plant biomass or forage production systems, WUE is equal to dry matter divided by evapotranspiration (Kramer and Kozlowski 1979). In agricultural production systems where grain yield is the product to be used or sold, WUE is equal to the ratio of grain yield to evapotranspiration (Viets 1962). However, evapotranspiration is difficult to measure, and therefore effective rainfall and irrigation are often used instead of evapotranspiration to calculate WUE in agronomic research (Howell 2001). In addition, depending on the source of water, some studies have also divided WUE into precipitation use efficiency (Peng *et al.* 2020) and irrigation use efficiency (Cao *et al.* 2020).

Reducing the input of water in agricultural production

usually causes a decrease in ET, and most crops suffer a reduction in yield when ET decreases (Evans and Sadler 2008; Nielsen *et al.* 2011). Therefore, the main way to improve water use efficiency is to improve water management capabilities. One method for increasing WUE is to change planting methods. No-tillage (Nielsen *et al.* 2005), furrow planting and straw mulching can increase WUE in arid areas (Peng *et al.* 2020). No-tillage can improve soil structure and crop root systems, and increase water absorption (Kan *et al.* 2020). Ditch planting and straw mulching can synchronize seasonal soil water supply with crop water needs (Wang *et al.* 2011). Another method for increasing WUE is to change the irrigation system. By using drip irrigation technology in the North China Plain, WUE can reach 2.27 kg m^{-3} , much higher than 1.45 kg m^{-3} observed when using surface irrigation (Zhang *et al.* 2011; Si *et al.* 2020). In areas where agricultural water is scarce, deficit irrigation can maximize the irrigation area, thereby achieving an increase in regional WUE (Bell *et al.* 2018; Li *et al.* 2019; Pardo *et al.* 2020). Additionally, changes in irrigation timing can affect WUE. Winter wheat is the most sensitive to moisture during jointing and flowering. Irrigation applications during these two growth stages can increase WUE by $0.45\text{--}0.96 \text{ kg m}^{-3}$ (Bai *et al.* 2020).

3.3. Radiation utilization efficiency

Radiation utilization efficiency (RdUE) is defined as the dry matter produced by solar radiation or photosynthetically active radiation per unit area intercepted by the canopy. It is an important quantitative indicator of crop yield related to photosynthesis (Stöckle and Kemanian 2009). Loomis and Amthor (1999) calculated the theoretical RdUE of C_3 crops as $1.5\text{--}2.0 \text{ g MJ}^{-1}$, and RdUE of C_4 crops as $4.0\text{--}5.8 \text{ g MJ}^{-1}$. However, the results of current research have shown that the actual RdUE of farmland crops is far lower than these theoretical values, and there are great differences in different regions. For example, the RdUE of maize in the United States can reach $3.74\text{--}3.84 \text{ g MJ}^{-1}$ (Lindquist *et al.* 2005), in China it was reported to be $0.90\text{--}1.69 \text{ g MJ}^{-1}$ (Jia *et al.* 2018), however, the RdUE in Mexico was only $0.54\text{--}0.68 \text{ g MJ}^{-1}$ (Morales Ruiz *et al.* 2016). RdUE is mainly influenced by crop genetics. Differences in RdUE between C_3 and C_4 plants due to different photosynthetic pathways are obvious. Rice is a C_3 crop that has been shown to have increased ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) regulation through genetic engineering technology in which introduced traits similar to those found in C_4 plants (such as CO_2 concentration mechanism) can improve light interception and canopy photosynthesis, and increase RdUE (Brar and Khush 2013; Brar *et al.* 2017). In addition, water and nitrogen supply, planting density, and

other agricultural management practices can also affect the RdUE of crops (Hatfield and Dold 2019).

4. Concluding remarks

Yield gap research has made many achievements in revealing the factors that limit crop yields, thereby improving farmers' yields. However, the following deficiencies in yield gap research exist:

(1) Lack of standardization. For the same variety in the same region, the potential yield and yield gap obtained in different studies can be very different. This is primarily a result of having no unified standard definition of potential yield, and researchers setting different yield levels corresponding to potential yield based on their specific needs. Additionally, because of differences in research methods, uncertainties in crop models, and the different emphasis of different models, there is a disagreement regarding the quantitative results representing potential yields from modeling.

(2) Poor regional accuracy. Both field trials and model simulations are based on site scale. When expanding the research results to the regional scale, differences in management methods (variety, fertilization, etc.) are usually ignored, but these factors directly affect crop yields.

(3) Incomplete research. Current research focuses on the time and space variability of yield gap, and the increase yield after closing yield gap (e.g., suppose it reaches 80% of the Y_p). However, almost none of them can accurately answer how to close the yield gap. Crop models cannot quantify the effects of yield-limiting factors; and the boundary line analysis method separates the interaction between factors, which is not in line with reality. Therefore, the exploration of yield-limiting factors and ways to close the yield gap almost only appeared in the discussion part of the paper, as some possibilities were proposed.

(4) Limited effect. The aim of the researchers is mostly to reduce the yield gap to ensure food security. So they pay more attention to the impact of crop management on yield, e.g., water and nitrogen supply. However, they ignore whether narrowing the yield gap can bring economic benefits to farmers, and whether RUE will decrease after changing crop management. Therefore, farmers are unwilling to follow the suggestions of researchers, which weakens the practical value of research to a certain extent.

Different from the yield gap research, RUE research focuses more on small scale. Except nutrient utilization efficiency, researches on water utilization efficiency and radiation utilization efficiency are mostly based on leaf and canopy scales. And RUE research usually only studies the utilization rate of a single resource. The problem with this approach is that an increase in RUE may be based

on the premise that the utilization rate of other resources decreases. For example, yield can be increased by increasing irrigation application amounts. If the input of fertilizers has not increased, then the efficiency of nitrogen use has indeed increased, but it is not clear whether the efficiency of water utilization has improved as well. In addition, there is little research regarding the potential RUE of farmland. What is the gap between the actual and the potential RUE? Knowing the answers to these questions will have great significance for guiding and determining actual production practices used by farmers.

Agricultural production is the result of the interaction of many factors. On the basis of understanding the temporal and spatial changes of the yield gap, comprehensive research on crop management, climate and socioeconomic factors, and quantifying the role of yield-limiting factors in the yield gap will help formulate measures to reduce the yield gap. Machine learning has advantages when dealing with multi-variables and complex data, it will be a good choice to use it to analyze the yield gap. At the same time, regional scale research should be combined with remote sensing of normalized vegetation index (NDVI), enhanced vegetation index (EVI) and other data to assist in correcting and improving simulation processes in crop models. Combining results from controlled field experiments with regional statistical data can also be used to improve the accuracy of regional model simulations. By dating new methods and data, rather than just using model simulations, it will help us make suggestions that are more acceptable to farmers to achieve the two goals of high yield and high efficiency.

Acknowledgements

This work was supported by the National Key Research and Development Program of China (2016YFD0300100), the Innovation Program of Chinese Academy of Agricultural Sciences and the Elite Youth Program of the Chinese Academy of Agricultural Science. We thank Prof. Yang Xiaoguang (China Agricultural University) for thoughtful comments on this manuscript.

Appendices associated with this paper can be available on <http://www.ChinaAgriSci.com/V2/En/appendix.htm>

References

- Agus F, Andrade J F, Edreira J I R, Deng N Y, Purwantomo D K G, Agustiani N, Aristya V E, Batubara S F, Herniwati, Hosang E Y, Krisnadi L Y, Makka A, Samijan, Cenacchi N, Wiebe K, Grassini P. 2019. Yield gaps in intensive rice–maize cropping sequences in the humid tropics of Indonesia. *Field Crops Research*, **237**, 12–22.
- Assefa B T, Chamberlin J, Reidsma P, Silva J V, van Ittersum M K. 2020. Unravelling the variability and causes of smallholder maize yield gaps in Ethiopia. *Food Security*, **12**, 83–103.
- Asseng S, Ewert F, Rosenzweig C, Jones J W, Hatfield J L, Ruane A C, Boote K J, Thorburn P J, Rötter R P, Cammarano D, Brisson N, Basso B, Martre P, Aggarwal P K, Angulo C, Bertuzzi P, Biernath C, Challinor A J, Doltra J, Gayler S, et al. 2013. Uncertainty in simulating wheat yields under climate change. *Nature Climate Change*, **3**, 827–832.
- Bai H Q, Wang J, Fang Q X, Huang B X. 2020. Does a trade-off between yield and efficiency reduce water and nitrogen inputs of winter wheat in the North China Plain? *Agricultural Water Management*, **233**, 106095.
- Balboa G R, Archontoulis S V, Salvagiotti F, Garcia F O, Stewart W M, Francisco E, Prasad P V V, Ciampitti I A. 2019. A systems-level yield gap assessment of maize–soybean rotation under high- and low-management inputs in the Western US Corn Belt using APSIM. *Agricultural Systems*, **174**, 145–154.
- Barker R K, Gomez A, Herdt R W. 1979. *Farm-level Constraints to High Rice Yields in Asia: 1974–77*. IRRI, Los Banos, Philippines.
- Bell J M, Schwartz R, McInnes K J, Howell T, Morgan C L S. 2018. Deficit irrigation effects on yield and yield components of grain sorghum. *Agricultural Water Management*, **203**, 289–296.
- de Bie C A J M. 2000. Comparative performance analysis of agro-ecosystems. Ph D thesis, Wageningen University, Wageningen, Netherlands.
- Bindraban P S, Stoorvogel J J, Jansen D M, Vlaming J, Groot J J R. 2000. Land quality indicators for sustainable land management: Proposed method for yield gap and soil nutrient balance. *Agriculture, Ecosystems & Environment*, **81**, 103–112.
- Brar D S, Khush G S. 2013. Chapter 8 — Biotechnological approaches for increasing productivity and sustainability of rice production. In: *Agricultural Sustainability*. Academic Press, San Diego, CA, USA. pp. 151–175.
- Brar D S, Singh K, Khush G S. 2017. Chapter 6 — Frontiers in rice breeding. In: *The Future Rice Strategy for India*. Academic Press, Cambridge, MA, USA. pp. 137–160.
- Bruinsma J. 2003. World agriculture: Towards 2015/2030: An FAO perspective. Earthscan, London and FAO, Rome.
- Cao H X, Zhao S L, Ge D K, Liu Y X, Liu Y, Sun J Y, Yue Y B, Zhang Z Y, Chen Y L. 2011. Discussion on development of crop models. *Scientia Agricultura Sinica*, **44**, 3520–3528. (in Chinese)
- Cao H Z, Li Y N, Chen G F, Chen D D, Qu H R, Ma W Q. 2019. Identifying the limiting factors driving the winter wheat yield gap on smallholder farms by agronomic diagnosis in North China Plain. *Journal of Integrative Agriculture*, **18**, 1701–1713.
- Cao X, Zeng W, Wu M, Guo X, Wang W. 2020. Hybrid analytical framework for regional agricultural water resource utilization and efficiency evaluation. *Agricultural Water Management*,

- 231, 106027.
- Carciochi W D, Sadras V O, Pagani A, Ciampitti I A. 2020. Co-limitation and stoichiometry capture the interacting effects of nitrogen and sulfur on maize yield and nutrient use efficiency. *European Journal of Agronomy*, **113**, 125973.
- Cassman K G, Dobermann A, Walters D T, Yang H. 2003. Meeting cereal demand while protecting natural resources and improving environmental quality. *Annual Review of Environment and Resources*, **28**, 315–358.
- Chen G, Cao H, Liang J, Ma W, Guo L, Zhang S, Jiang R, Zhang H, Goulding K W, Zhang F. 2018. Factors affecting nitrogen use efficiency and grain yield of summer maize on smallholder farms in the North China Plain. *Sustainability*, **10**, 363.
- Cheng M, Wang R H, Xue H X, Li Q. 2012. Effects of drought on ecosystem net primary production in northwestern China. *Journal of Arid Land Resources and Environment*, **6**, 1–7. (in Chinese)
- Cirera X, Masset E. 2010. Income distribution trends and future food demand. *Philosophical Transactions of the Royal Society (B: Biological Sciences)*, **365**, 2821–2834.
- Cui Z, Wang G, Yue S, Wu L, Zhang W, Zhang F, Chen X. 2014. Closing the N-use efficiency gap to achieve food and environmental security. *Environmental Science & Technology*, **48**, 5780–5787.
- Dai X Q, Ouyang Z, Li Y S, Wang H M. 2013. Variation in yield gap induced by nitrogen, phosphorus and potassium fertilizer in North China Plain. *PLoS ONE*, **8**, e82147.
- de Datta S K. 1981. *Principles and Practices of Rice Production*. Wiley-Interscience Productions, New York, USA.
- Ding Y M, Wang W G, Zhuang Q L, Luo Y F. 2020. Adaptation of paddy rice in China to climate change: The effects of shifting sowing date on yield and irrigation water requirement. *Agricultural Water Management*, **228**, 105890.
- Dutta S, Chakraborty S, Goswami R, Banerjee H, Majumdar K, Li B, Jat M L. 2020. Maize yield in smallholder agriculture system — An approach integrating socio-economic and crop management factors. *PLoS ONE*, **15**, e0229100.
- Espe M B, Cassman K G, Yang H, Guilpart N, Grassini P, Van Wart J, Anders M, Beighley D, Harrell D, Linscombe S, McKenzie K, Mutters R, Wilson L T, Linquist B A. 2016. Yield gap analysis of US rice production systems shows opportunities for improvement. *Field Crops Research*, **196**, 276–283.
- Espe M B, Hill J E, Leinfelder-Miles M, Espino L A, Mutters R, Mackill D, van Kessel C, Linquist B A. 2018. Rice yield improvements through plant breeding are offset by inherent yield declines over time. *Field Crops Research*, **222**, 59–65.
- Evans R G, Sadler E J. 2008. Methods and technologies to improve efficiency of water use. *Water Resources Research*, **44**, 1–15.
- Evenson R E, Gollin D. 2003. Assessing the impact of the green revolution, 1960 to 2000. *Science*, **300**, 758–762.
- Ewert F, Porter J R, Rounsevell M D A. 2007. Crop models, CO₂, and climate change. *Science*, **315**, 459–460.
- Fan M, Shen J, Yuan L, Jiang R, Chen X, Davies W J, Zhang F. 2011. Improving crop productivity and resource use efficiency to ensure food security and environmental quality in China. *Journal of Experimental Botany*, **63**, 13–24.
- Fan S, Wailes E J, Young K B. 1997. Policy reforms and technological change in Egyptian rice production: A frontier production function approach. *Journal of African Economies*, **6**, 391–411.
- Fang G, Wen Y, Yu H, van der Werf W, Qiang C, Heerink N, van Ittersum M K. 2017. On yield gaps and yield gains in intercropping: Opportunities for increasing grain production in northwest China. *Agricultural Systems*, **151**, 96–105.
- FAO (Food and Agriculture Organization). 2009. *How to Feed the World in 2050*. High Level Expert Forum, Rome.
- Fermont A M, van Asten P J A, Titttonell P, van Wijk M T, Giller K E. 2009. Closing the cassava yield gap: An analysis from smallholder farms in East Africa. *Field Crops Research*, **112**, 24–36.
- Fitzgerald T, Kuwayama Y, Olmstead S, Thompson A. 2020. Dynamic impacts of U.S. energy development on agricultural land use. *Energy Policy*, **137**, 111163.
- Fixen P, Brentrup F, Bruulsema T, Garcia F, Norton R, Zingore S. 2015. Chapter 8 — Nutrient/Fertilizer use efficiency: measurement, current situation and trends. In: *Managing Water and Fertilizer for Sustainable Agricultural Intensification*. International Fertilizer Industry Association, Paris, France. pp. 8–37.
- Fresco L. 1984. Issues in farming systems research. *Netherlands Journal of Agricultural Science*, **32**, 253–261.
- Gibbs H K, Ruesch A S, Achard F, Clayton M K, Holmgren P, Ramankutty N, Foley J A. 2010. Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 16732–16737.
- Hajjarpoor A, Soltani A, Zeinali E, Kashiri H, Ayneband A, Vadez V. 2018. Using boundary line analysis to assess the on-farm crop yield gap of wheat. *Field Crops Research*, **225**, 64–73.
- Hardke J, Schmidt L, Mazzanti R. 2013. 2013 Arkansas rice quick facts. University of Arkansas Division of Agriculture, Research and Extension. [2013-08-08]. http://www.aragriculture.org/crops/rice/quick_facts/2013_rice_quick_facts.pdf
- Hatfield J L, Boote K J, Kimball B A, Ziska L H, Izaurralde R C, Ort D, Thomson A M, Wolfe D. 2011. Climate impacts on agriculture: Implications for crop production. *Agronomy Journal*, **103**, 351–370.
- Hatfield J L, Dold C. 2019. Chapter 1 — Photosynthesis in the solar corridor system. In: *The Solar Corridor Crop System*. Academic Press, Cambridge, MA, USA. pp. 1–33.
- Hochman Z, Gobbett D, Holzworth D, McClelland T, van Rees H, Marinoni O, Garcia J N, Horan H. 2013. Reprint of “Quantifying yield gaps in rainfed cropping systems: A case study of wheat in Australia”. *Field Crops Research*, **143**, 65–75.
- Hoffmann M P, Haakana M, Asseng S, Höhn J G, Palosuo T, Ruiz-Ramos M, Fronzek S, Ewert F, Gaiser T, Kassie B T,

- Paff K, Rezaei E E, Rodríguez A, Semenov M, Srivastava A K, Stratonovitch P, Tao F, Chen Y, Rötter R P. 2018. How does inter-annual variability of attainable yield affect the magnitude of yield gaps for wheat and maize? An analysis at ten sites. *Agricultural Systems*, **159**, 199–208.
- Howell T A. 2001. Enhancing water use efficiency in irrigated agriculture. *Agronomy Journal*, **93**, 281–289.
- Hu X, Huang Y, Sun W, Yu L. 2017. Shifts in cultivar and planting date have regulated rice growth duration under climate warming in China since the early 1980s. *Agricultural and Forest Meteorology*, **247**, 34–41.
- van Ittersum M K, Cassman K G, Grassini P, Wolf J, Tittonell P, Hochman Z. 2013. Yield gap analysis with local to global relevance — A review. *Field Crops Research*, **143**, 4–17.
- Jia Q M, Sun L F, Mou H Y, Ali S, Liu D H, Zhang Y, Zhang P, Ren X L, Jia Z K. 2018. Effects of planting patterns and sowing densities on grain-filling, radiation use efficiency and yield of maize (*Zea mays* L.) in semi-arid regions. *Agricultural Water Management*, **201**, 287–298.
- Jin J Y. 2012. Changes in the efficiency of fertiliser use in China. *Journal of the Science of Food and Agriculture*, **92**, 1006–1015.
- Kan Z R, Liu Q Y, He C, Jing Z H, Virk A L, Qi J Y, Zhao X, Zhang H L. 2020. Responses of grain yield and water use efficiency of winter wheat to tillage in the North China Plain. *Field Crops Research*, **249**, 107760.
- Khaliq T, Gaydon D S, Ahmad M U D, Cheema M J M, Gull U. 2019. Analyzing crop yield gaps and their causes using cropping systems modelling — A case study of the Punjab rice–wheat system, Pakistan. *Field Crops Research*, **232**, 119–130.
- Kramer P J, Kozlowski T T. 1979. *Physiology of Woody Plants*. Academic Press, New York.
- Kukul M S, Irmak S. 2018. Climate-driven crop yield and yield variability and climate change impacts on the U.S. great plains agricultural production. *Scientific Reports*, **8**, 3450.
- Laborte A G, de Bie K, Smaling E M A, Moya P F, Boling A A, Van Ittersum M K. 2012. Rice yields and yield gaps in Southeast Asia: Past trends and future outlook. *European Journal of Agronomy*, **36**, 9–20.
- Li K N, Yang X G, Liu Z J, Zhang T Y, Lu S, Liu Y. 2014. Low yield gap of winter wheat in the North China Plain. *European Journal of Agronomy*, **59**, 1–12.
- Li Q, Cui X, Liu X, Roelcke M, Pasda G, Zerulla W, Wissemeyer A H, Chen X, Goulding K, Zhang F. 2017. A new urease-inhibiting formulation decreases ammonia volatilization and improves maize nitrogen utilization in North China Plain. *Scientific Reports*, **7**, 43853.
- Li X D, Du Y, Wu S J, Sun J Y, Feng Q, Song Y J. 2009. Evolution and effects of climate-productivity on climate change in Hubei Province. *System Sciences and Comprehensive Studies in Agriculture*, **25**, 294–298. (in Chinese).
- Li X M, Zhao W X, Li J S, Li Y F. 2019. Maximizing water productivity of winter wheat by managing zones of variable rate irrigation at different deficit levels. *Agricultural Water Management*, **216**, 153–163.
- Licker R, Johnston M, Foley J A, Barford C, Kucharik C J, Monfreda C, Ramankutty N. 2010. Mind the gap: How do climate and agricultural management explain the ‘yield gap’ of croplands around the world? *Global Ecology and Biogeography*, **19**, 769–782.
- Lin Z H, Mo X G, Xiang Y Q. 2003. Research advances on crop growth models. *Acta Agronomica Sinica*, **5**, 750–758. (in Chinese)
- Lindquist J L, Arkebauer T J, Walters D T, Cassman K G, Dobermann A. 2005. Maize radiation use efficiency under optimal growth conditions. *Agronomy Journal*, **97**, 72–78.
- Linquist B, Snyder R, Anderson F, Espino L, Inglese G, Marras S, Moratiel R, Mutters R, Nicolosi P, Rejmanek H, Russo A, Shapland T, Song Z, Swelam A, Tindula G, Hill J. 2015. Water balances and evapotranspiration in water- and dry-seeded rice systems. *Irrigation Science*, **33**, 375–385.
- Liu B H, Chen X P, Meng Q F, Yang H S, van Wart J. 2017. Estimating maize yield potential and yield gap with agro-climatic zones in China — Distinguish irrigated and rainfed conditions. *Agricultural and Forest Meteorology*, **239**, 108–117.
- Liu J D, Zhou X J, Yu Q. 2001. Modification of the basic parameters in FAO productivity model. *Journal of Natural Resources*, **16**, 240–247. (in Chinese)
- Liu W, Lv P, Su K, Yang J S, Zhang J W, Dong S T, Liu P, Sun Q Q. 2010. Effects of planting density on the grain yield and source-sink characteristics of summer maize. *Journal of Applied Ecology*, **21**, 1737–1743. (in Chinese)
- Liu Y, Wang E, Yang X, Wang J. 2010. Contributions of climatic and crop varietal changes to crop production in the North China Plain, since 1980s. *Global Change Biology*, **16**, 2287–2299.
- Liu Z J, Yang X G, Hubbard K G, Lin X M. 2012. Maize potential yields and yield gaps in the changing climate of northeast China. *Global Change Biology*, **18**, 3441–3454.
- Lobell D B, Cassman K G, Field C B. 2009. Crop yield gaps: Their importance, magnitudes, and causes. *Annual Review of Environment and Resources*, **34**, 179–204.
- Lobell D B, Ortiz-Monasterio J I. 2006. Regional importance of crop yield constraints: Linking simulation models and geostatistics to interpret spatial patterns. *Ecological Modelling*, **196**, 173–182.
- Long S P, Ainsworth E A, Leakey A D B, Nösberger J, Ort D R. 2006. Food for thought: Lower-than-expected crop yield stimulation with rising CO₂ concentrations. *Science*, **312**, 1918–1921.
- Loomis R S, Amthor J. 1999. Yield potential, plant assimilatory capacity, and metabolic efficiencies. *Crop Science*, **39**, 1584–1596.
- van Loon M P, Adjei-Nsiah S, Descheemaeker K, Akotsen-Mensah C, van Dijk M, Morley T, van Ittersum M K, Reidsma P. 2019. Can yield variability be explained? Integrated assessment of maize yield gaps across smallholders in Ghana. *Field Crops Research*, **236**, 132–144.
- Lu C, Li X, Tan M. 2007. Chapter 3 — China’s farmland use: A

- scenario analysis of changes and trends. In: *Dragons with Clay Feet? Transition, Sustainable Rural Resource Use, and Rural Environment in China and Vietnam*. Rowman and Littlefield Publishers, Lanham and Boston, USA. pp. 309–326.
- Lu C H, Fan L. 2013. Winter wheat yield potentials and yield gaps in the North China Plain. *Field Crops Research*, **143**, 98–105.
- Luo Y Z, Cheng Z Y, Guo X Q. 2011. The changing characteristics of potential climate productivity in Gansu Province during nearly 40 years. *Shengtai Xuebao/Acta Ecologica Sinica*, **31**, 221–229. (in Chinese)
- Maas S J. 1988. Use of remotely-sensed information in agricultural crop growth models. *Ecological Modelling*, **41**, 247–268.
- Mahmood N, Arshad M, Kächele H, Ma H, Ullah A, Müller K. 2019. Wheat yield response to input and socioeconomic factors under changing climate: Evidence from rainfed environments of Pakistan. *Science of The Total Environment*, **688**, 1275–1285.
- Mao Y D, Su G L, Li F D, Wu Y X. 2008. Impact of climate change on plant climate productivity in Zhejiang Province. *Chinese Journal of Eco-agriculture*, **16**, 273–278. (in Chinese)
- Maria Carolina D S A, Boote K J, Sentelhas P C, Romanelli T L. 2018. Variability and limitations of maize production in Brazil: Potential yield, water-limited yield and yield gaps. *Agricultural Systems*, **165**, 264–273.
- Meertens H, Ndege L, Enserink H. 1991. Results of the urea demonstration on-farm trial: Maswa District 1990–91. Field Note No. 26. Tanzania/Netherlands Farming System Research Project, Lake Zone, Mwanza, Tanzania.
- Meng Q, Yue S, Hou P, Cui Z, Chen X. 2016. Improving yield and nitrogen use efficiency simultaneously for maize and wheat in China: A review. *Pedosphere*, **26**, 137–147.
- Morales-Ruiz A, Loeza-Corte J, Díaz-López E, Morales-Rosales E, Franco-Mora O, Mariezcurrena-Berasain M, Estrada-Campuzano G. 2016. Efficiency on the use of radiation and corn yield under three densities of sowing. *International Journal of Agronomy*, **2016**, 1–5.
- Moulin S, Bondeau A, Delecalle R. 1998. Combining agricultural crop models and satellite observations: From field to regional scales. *International Journal of Remote Sensing*, **19**, 1021–1036.
- Mueller N D, Gerber J S, Johnston M, Ray D K, Ramankutty N, Foley J A. 2012. Closing yield gaps through nutrient and water management. *Nature*, **490**, 254–257.
- Nalley L, Tack J, Barkley A, Jagadish K, Brye K. 2016. Quantifying the agronomic and economic performance of hybrid and conventional rice varieties. *Agronomy Journal*, **108**, 1514–1523.
- Nielsen D C, Unger P W, Miller P R. 2005. Efficient water use in dryland cropping systems in the Great Plains. *Agronomy Journal*, **97**, 364–372.
- Nielsen D C, Vigil M F, Benjamin J G. 2011. Evaluating decision rules for dryland rotation crop selection. *Field Crops Research*, **120**, 254–261.
- van Oort P A J, Saito K, Dieng I, Grassini P, Cassman K G, van Ittersum M K. 2017. Can yield gap analysis be used to inform R&D prioritisation? *Global Food Security*, **12**, 109–118.
- Pardey P G, Beddow J M, Hurley T M, Beatty T K, Eidman V R. 2014. A bounds analysis of world food futures: Global agriculture through to 2050. *Australian Journal of Agricultural and Resource Economics*, **58**, 571–589.
- Pardo J J, Martínez-Romero A, Lélis B C, Tarjuelo J M, Domínguez A. 2020. Effect of the optimized regulated deficit irrigation methodology on water use in barley under semiarid conditions. *Agricultural Water Management*, **228**, 105925.
- Patrignani A, Lollato R P, Ochsner T E, Godsey C B, Edwards J T. 2014. Yield gap and production gap of rainfed winter wheat in the Southern Great Plains. *Agronomy Journal*, **106**, 1329–1339.
- Pellegrini P, Fernández R J. 2018. Crop intensification, land use, and on-farm energy-use efficiency during the worldwide spread of the green revolution. *Proceedings of the National Academy of Sciences the United States of America*, **115**, 2335–2340.
- Peng S, Khush G S, Virk P, Tang Q, Zou Y. 2008. Progress in ideotype breeding to increase rice yield potential. *Field Crops Research*, **108**, 32–38.
- Peng Z K, Wang L L, Xie J H, Li L L, Coulter J A, Zhang R Z, Luo Z Z, Cai L Q, Carberry P, Whitbread A. 2020. Conservation tillage increases yield and precipitation use efficiency of wheat on the semi-arid Loess Plateau of China. *Agricultural Water Management*, **231**, 106024.
- Rabbinge R. 1993. The ecological background of food production. In: Chadwick D J, Marsh J, eds., *Crop Protection and Sustainable Agriculture (Ciba Foundation Symposium 177)*. Wiley, Chichester, UK. pp. 2–29.
- Ramankutty N, Foley J A, Norman J, McSweeney K. 2002. The global distribution of cultivable lands: Current patterns and sensitivity to possible climate change. *Global Ecology and Biogeography*, **11**, 377–392.
- Rosegrant M W, Paisner M S, Meijer S, Witcover J. 2001. *Global Food Projections to 2020: Emerging Trends and Alternative Futures*. International Food Policy Research Institute, Washington, USA.
- Sadras V O. 2006. The N:P stoichiometry of cereal, grain legume and oilseed crops. *Field Crops Research*, **95**, 13–29.
- Sarobol N, Virakul P, Potan N, Benjasil V, Setarath P, Dechates S. 1989. *Preliminary Survey on Soybean Yield Gap Analysis in Thailand*. The CGPRT Centre Publication, Bogor, Indonesia.
- Senapati N, Semenov M A. 2020. Large genetic yield potential and genetic yield gap estimated for wheat in Europe. *Global Food Security*, **24**, 100340.
- Shen J B, Cui Z L, Miao Y X, Mi G H, Zhang H Y, Fan M S, Zhang C C, Jiang R F, Zhang W F, Li H G, Chen X P, Li X L, Zhang F S. 2013. Transforming agriculture in China: From solely high yield to both high yield and high resource use efficiency. *Global Food Security*, **2**, 1–8.

- Si Z Y, Zain M, Mehmood F, Wang G S, Gao Y, Duan A. 2020. Effects of nitrogen application rate and irrigation regime on growth, yield, and water-nitrogen use efficiency of drip-irrigated winter wheat in the North China Plain. *Agricultural Water Management*, **231**, 106002.
- Silva J V, Reidsma P, Laborte A G, van Ittersum M K. 2017. Explaining rice yields and yield gaps in Central Luzon, Philippines: An application of stochastic frontier analysis and crop modelling. *European Journal of Agronomy*, **82**, 223–241.
- Stöckle C O, Kemanian A R. 2009. Chapter 7 — Crop radiation capture and use efficiency: A framework for crop growth analysis. In: Sadras V, Calderini D, eds., *Crop Physiology*. Academic Press, San Diego.
- Takai T, Matsuura S, Nishio T, Ohsumi A, Shiraiwa T, Horie T. 2006. Rice yield potential is closely related to crop growth rate during late reproductive period. *Field Crops Research*, **96**, 328–335.
- Tao F, Yokozawa M, Xu Y, Hayashi Y, Zhang Z. 2006. Climate changes and trends in phenology and yields of field crops in China, 1981–2000. *Agricultural and Forest Meteorology*, **138**, 82–92.
- UNDESAPD (United Nations, Department of Economic and Social Affairs, Population Division). 2019. *World Population Prospects 2019: Highlights*. ST/ESA/SER.A/423.
- Valin H, Sands R D, Van der Mensbrugge D, Nelson G C, Ahammad H, Blanc E, Bodirsky B, Fujimori S, Hasegawa T, Havlik P. 2014. The future of food demand: Understanding differences in global economic models. *Agricultural Economics*, **45**, 51–67.
- Viets F G. 1962. Fertilizers and the efficient use of water. *Advances in Agronomy*, **14**, 223–264.
- Wang C, Li X L, Gong T T, Zhang H Y. 2014. Life cycle assessment of wheat-maize rotation system emphasizing high crop yield and high resource use efficiency in Quzhou County. *Journal of Cleaner Production*, **68**, 56–63.
- Wang J W, Zhang J H, Bai Y, Zhang S, Yang S S, Yao F M. 2020. Integrating remote sensing-based process model with environmental zonation scheme to estimate rice yield gap in Northeast China. *Field Crops Research*, **246**, 107682.
- Wang T C, Wei L, Wang H Z, Ma S C, Ma B L. 2011. Responses of rainwater conservation, precipitation-use efficiency and grain yield of summer maize to a furrow-planting and straw-mulching system in northern China. *Field Crops Research*, **124**, 223–230.
- Wilbois K P, Schmidt J E. 2019. Reframing the debate surrounding the yield gap between organic and conventional farming. *Agronomy*, **9**, 82.
- Wu D, Qiang Y, He L C, Hengsdijk H. 2006. Quantifying production potentials of winter wheat in the North China Plain. *European Journal of Agronomy*, **24**, 226–235.
- Xu X P, He P, Pampolino M F, Li Y Y, Liu S Q, Xie J G, Hou Y P, Zhou W. 2016. Narrowing yield gaps and increasing nutrient use efficiencies using the Nutrient Expert system for maize in Northeast China. *Field Crops Research*, **194**, 75–82.
- Xu Z, Li C, Zhang C, Yu Y, van der Werf W, Zhang F. 2020. Intercropping maize and soybean increases efficiency of land and fertilizer nitrogen use; A meta-analysis. *Field Crops Research*, **246**, 107661.
- Yu H L, Huang J Y, Wang T H. 2012. Impact of climate change on climate potential productivity in recent 56 years in middle arid regions of Ningxia — A case study of Zhongning County. *Research of Soil and Water Conservation*, **6**, 35. (in Chinese)
- Zeleele K T, Nendel C. 2016. Analysis of options for increasing wheat (*Triticum aestivum* L.) yield in south-eastern Australia: The role of irrigation, cultivar choice and time of sowing. *Agricultural Water Management*, **166**, 139–148.
- Zhang H, Hou D P, Peng X L, Ma B J, Shao S M, Jing W J, Gu J F, Liu L J, Wang Z Q, Liu Y Y, Yang J C. 2019a. Optimizing integrative cultivation management improves grain quality while increasing yield and nitrogen use efficiency in rice. *Journal of Integrative Agriculture*, **18**, 2716–2731.
- Zhang H, Tao F L, Zhou G S. 2019b. Potential yields, yield gaps, and optimal agronomic management practices for rice production systems in different regions of China. *Agricultural Systems*, **171**, 100–112.
- Zhang W, Cao G, Li X, Zhang H, Wang C, Liu Q, Chen X, Cui Z, Shen J, Jiang R, Mi G, Miao Y, Zhang F, Dou Z. 2016. Closing yield gaps in China by empowering smallholder farmers. *Nature*, **537**, 671–674.
- Zhang X Y, Chen S Y, Sun H Y, Shao L W, Wang Y Z. 2011. Changes in evapotranspiration over irrigated winter wheat and maize in North China Plain over three decades. *Agricultural Water Management*, **98**, 1097–1104.